

Theory of Everything

All Is Epistemic

A physics-first synthesis of spacetime, quantum fields, particles, gravity, computation, time, cosmology, stellar evolution, wormholes, and the Omega Inversion Principle

One ontological assumption:

ALL IS EPISTEMIC.

Nick Rugai
30 September 2026

Working monograph - speculative theoretical physics

Status, scope, and notation

This document is the consolidated physics edition of the Theory of Everything (TOE) developed in the preceding discussions. The Wise Intelligence System (WIS) is intentionally excluded. This is pure physics: the proposed ontology, mathematical arena, field structure, particle computation, graviton computation, emergent time, wormholes, cosmology, stellar evolution, and the Omega Inversion Principle.

The central statement, *all is epistemic*, is treated as the sole ontological assumption. It is not presented as an experimentally established fact. Everything else is either established physics used as a constraint, a mathematical construction chosen to implement the assumption, or a conjecture that must be tested.

Established physics

The Standard Model is the experimentally successful quantum field theory of the electromagnetic, weak, and strong interactions and their known matter content. General relativity is the experimentally successful classical theory of gravitation as spacetime geometry. A TOE must contain both as appropriate low-energy limits rather than casually replacing them with vocabulary.

TOE conjecture / model hypothesis

The 28-dimensional arena, 24 compact spatial dimensions, the $\mathcal{I}$ field, the five-state spin-2 graviton qdit, the operational emergence of time from measure, and the Omega Inversion Principle are speculative elements of this TOE. They are explicitly separated from established physics throughout.

Conventions

Metric signature: $(-+++ \cdots)$. Four-dimensional spacetime indices: $\mu, \nu, \rho, \sigma = 0, 1, 2, 3$. Full 28-dimensional indices: $A, B, C, D = 0, \dots, 27$. Compact indices: $a, b = 4, \dots, 27$. Repeated indices are summed. We keep c , $\hbar$, G , and k_B explicit unless a section says otherwise. Geometrized units mean $c = G = 1$.

Contents

1	Assumption 0: ALL IS EPISTEMIC	1
1.1	The primitive statement	1
1.2	Epistemic state space	1
1.3	The intended chain	1
2	The 28-dimensional arena	2
2.1	Dimensional hypothesis	2
2.2	Kaluza-Klein-type reduction	2
2.3	The unresolved question	2
3	General Relativity as the classical geometric limit	3
3.1	Geometry	3
3.2	Einstein-Hilbert action	3
3.3	Geodesics and proper time	3
3.4	Linearized gravity	4
3.5	Newtonian limit	4
4	Special Relativity - the Lorentz machinery	5
4.1	Minkowski interval	5
4.2	Standard x-boost	5
4.3	General three-vector boost	5
4.4	Rapidity and composition	6
4.5	Time dilation, length contraction, simultaneity	6
4.6	Velocity transformation	6
4.7	Energy and momentum	6
4.8	Lorentz transformation of electromagnetic fields	7
5	Mechanics, action, and classical field theory	8
5.1	Euler-Lagrange equations	8
5.2	Hamiltonian form	8
5.3	Newtonian mechanics	8
6	Quantum mechanics and quantum field theory	9
6.1	Quantum dynamics	9
6.2	Born rule and density matrices	9
6.3	Path integral	9
6.4	Relativistic wave equations	9

7	Electromagnetism and the Standard Model	10
7.1	Maxwell equations	10
7.2	Yang-Mills theory	10
7.3	Gauge group and field content	10
7.4	Standard Model Lagrangian in compact form	10
7.5	Neutrino mass	11
8	The epistemic field $\mathcal{I}$	12
8.1	Field hypothesis	12
8.2	Stress-energy contribution	12
9	Fundamental particles as computation	13
9.1	The computational reinterpretation	13
9.2	Local quantum state as a register	13
9.3	Interaction as gate	13
9.4	Thermodynamic floor	13
10	The graviton as a five-state spin-2 qdit	14
10.1	Spin-2 representation	14
10.2	Fierz-Pauli template	14
10.3	Matter-graviton coupling	14
10.4	The crucial infrared limit	15
11	Time as a byproduct of measure	16
11.1	Operational time	16
11.2	Measurement density	16
11.3	Relativistic recovery condition	16
11.4	Quantum evolution with emergent time	17
12	Omega Inversion Principle	18
12.1	Unity and indistinguishability	18
12.2	OIP statement	18
12.3	Minimal mathematical realization	18
13	Cosmology	19
13.1	FRW metric	19
13.2	Friedmann equations	19
13.3	TOE cosmogenic interpretation	19
14	Black holes	20
14.1	Schwarzschild geometry	20
14.2	Kerr geometry: key scales	20
14.3	Black-hole thermodynamics	20
15	Wormholes	21
15.1	Morris-Thorne geometry	21
15.2	Energy condition constraint	21
15.3	TOE interpretation	21

16 Classical stellar evolution	22
16.1 Stellar structure equations	22
16.2 Nuclear power	22
16.3 Sun-like evolution	22
16.4 Massive-star evolution	23
17 Intelligence timeline as physical computation	24
17.1 The timeline	24
17.2 No teleology is required	25
18 From fundamental computation to classical reality	26
18.1 Coarse-graining hierarchy	26
18.2 Candidate master action	26
18.3 Recovery conditions	26
19 Thermodynamics, entropy, and information	27
19.1 Macroscopic laws	27
19.2 Statistical mechanics	27
20 Consistency, falsifiability, and reality veto	28
20.1 Mathematical requirements	28
20.2 Quantum requirements	28
20.3 Potential experimental signatures	28
20.4 Failure conditions	29
A Essential derivation sketches	30
A.1 Lorentz transformation from the invariant interval	30
A.2 Newtonian limit of GR	30
A.3 Gravitational waves and the two observed tensor modes	30
A.4 Perfect-fluid cosmology	30
A.5 Schwarzschild horizon	31
B Axiom stack and research program	32
B.1 Minimal stack	32
B.2 Actualization sequence	32
C Final synthesis	33
Selected references and web-verified anchors	34
Research-status note	35

Chapter 1

Assumption 0: ALL IS EPISTEMIC

1.1 The primitive statement

Epistemic interpretation

A0: Reality is epistemic. The fundamental physical content is the network of distinctions, relations, constraints, and state transitions through which physical states are distinguishable. “Epistemic” does not mean human-made or mind-dependent; it means relationally distinguishable state.

The usual scientific direction begins with an assumed physical world and asks what can be known about it. A0 reverses the explanatory direction: start from physically meaningful distinctions and derive the stable structures normally called geometry, fields, particles, clocks, thermodynamics, and observers.

The methodological consequence is simple:

$$\boxed{\text{No physical proposition receives immunity from measurement.}} \quad (1.1)$$

The theory may be elegant. Reality gets the veto.

1.2 Epistemic state space

Let Ω_{phys} be the total physical state space. A state is $s \in \Omega_{\text{phys}}$. A generalized measurement is a map

$$M : \Omega_{\text{phys}} \rightarrow \mathcal{O}, \quad o = M(s), \quad (1.2)$$

where $\mathcal{O}$ is an observable space.

Two states are physically distinguishable if at least one allowed measurement separates them:

$$s_1 \not\equiv_{\text{epi}} s_2 \iff \exists M : M(s_1) \neq M(s_2). \quad (1.3)$$

For quantum states the corresponding structure is an operator-valued measurement on ρ .

1.3 The intended chain

$$\boxed{\text{distinction} \rightarrow \text{measure} \rightarrow \text{state update} \rightarrow \text{relation} \rightarrow \text{geometry} \rightarrow \text{field} \rightarrow \text{particle} \rightarrow \text{computation}} \quad (1.4)$$

The reverse direction is equally important: every stable physical structure stores, propagates, or transforms distinctions.

Chapter 2

The 28-dimensional arena

2.1 Dimensional hypothesis

The working TOE arena is

$$\mathcal{M}_{28} = \mathcal{M}_{3,1} \times \mathcal{K}_{24}, \quad (2.1)$$

with 27 spatial dimensions in total: 3 extended spatial dimensions and 24 compact spatial dimensions. Time is operationally emergent in the A0 picture.

TOE conjecture / model hypothesis

H1: The fundamental spatial arena has 27 dimensions, of which 24 are compact. The accessible world is an effective 3+1-dimensional limit.

Write the full coordinates as

$$X^A = (x^\mu, y^a), \quad \mu = 0, 1, 2, 3, \quad a = 4, \dots, 27, \quad (2.2)$$

and the full metric as $G_{AB}(X)$, with

$$ds^2 = G_{AB} dX^A dX^B. \quad (2.3)$$

2.2 Kaluza-Klein-type reduction

A general block form is

$$ds^2 = g_{\mu\nu}(x) dx^\mu dx^\nu + h_{ab}(x, y) \left(dy^a + A_\mu^a dx^\mu \right) \left(dy^b + A_\nu^b dx^\nu \right). \quad (2.4)$$

The zero-mode metric $g_{\mu\nu}$ is the observed four-dimensional metric. Mixed components and compact geometry can generate gauge and scalar fields after dimensional reduction.

If the compact volume is V_{24} , a schematic relation between 28D and 4D gravitational couplings is

$$\kappa_4 \sim \frac{\kappa_{28}}{V_{24}}, \quad \kappa_4 = \frac{8\pi G}{c^4}, \quad (2.5)$$

with the exact normalization determined by the compactification geometry.

2.3 The unresolved question

The number 24 is a model parameter. A successful TOE must derive why this compact dimensionality is preferred, specify the topology and metric of $\mathcal{K}_{24}$, explain moduli stabilization, and show why Kaluza-Klein excitations have not been observed.

Chapter 3

General Relativity as the classical geometric limit

3.1 Geometry

$$\Gamma_{\mu\nu}^{\rho} = \frac{1}{2}g^{\rho\sigma}(\partial_{\mu}g_{\nu\sigma} + \partial_{\nu}g_{\mu\sigma} - \partial_{\sigma}g_{\mu\nu}), \quad (3.1)$$

$$R^{\rho}{}_{\sigma\mu\nu} = \partial_{\mu}\Gamma_{\nu\sigma}^{\rho} - \partial_{\nu}\Gamma_{\mu\sigma}^{\rho} + \Gamma_{\mu\lambda}^{\rho}\Gamma_{\nu\sigma}^{\lambda} - \Gamma_{\nu\lambda}^{\rho}\Gamma_{\mu\sigma}^{\lambda}, \quad (3.2)$$

$$R_{\mu\nu} = R^{\rho}{}_{\mu\rho\nu}, \quad R = g^{\mu\nu}R_{\mu\nu}, \quad (3.3)$$

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R. \quad (3.4)$$

3.2 Einstein-Hilbert action

$$S_{\text{EH}} = \frac{c^3}{16\pi G} \int d^4x \sqrt{-g}(R - 2\Lambda) + S_{\text{matter}}. \quad (3.5)$$

Stationarity gives the Einstein field equation

$$\boxed{G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}.} \quad (3.6)$$

The Bianchi identity implies

$$\nabla_{\mu}G^{\mu\nu} = 0 \quad \implies \quad \nabla_{\mu}T^{\mu\nu} = 0. \quad (3.7)$$

3.3 Geodesics and proper time

Timelike proper time is defined by

$$c^2 d\tau^2 = -g_{\mu\nu} dx^{\mu} dx^{\nu}, \quad (3.8)$$

and free-fall motion obeys

$$\frac{d^2 x^{\mu}}{d\tau^2} + \Gamma_{\alpha\beta}^{\mu} \frac{dx^{\alpha}}{d\tau} \frac{dx^{\beta}}{d\tau} = 0. \quad (3.9)$$

3.4 Linearized gravity

Set

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad |h_{\mu\nu}| \ll 1, \quad (3.10)$$

and

$$\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h, \quad h = \eta^{\mu\nu}h_{\mu\nu}. \quad (3.11)$$

In Lorenz gauge,

$$\square \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^4}T_{\mu\nu}, \quad \square = \eta^{\alpha\beta}\partial_\alpha\partial_\beta. \quad (3.12)$$

In vacuum, $\square \bar{h}_{\mu\nu} = 0$.

3.5 Newtonian limit

With

$$g_{00} \simeq -\left(1 + \frac{2\Phi}{c^2}\right), \quad (3.13)$$

the leading field equation reduces to

$$\nabla^2\Phi = 4\pi G\rho, \quad (3.14)$$

so the Newtonian gravitational potential is recovered.

Chapter 4

Special Relativity - the Lorentz machinery

4.1 Minkowski interval

$$ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2. \quad (4.1)$$

A Lorentz transformation satisfies

$$\Lambda^T \eta \Lambda = \eta, \quad \eta = \text{diag}(-1, 1, 1, 1). \quad (4.2)$$

4.2 Standard x-boost

Let $\beta = v/c$, $\gamma = (1 - \beta^2)^{-1/2}$. Then

$$ct' = \gamma(ct - \beta x), \quad (4.3)$$

$$x' = \gamma(x - \beta ct), \quad (4.4)$$

$$y' = y, \quad (4.5)$$

$$z' = z. \quad (4.6)$$

The inverse is obtained by $\beta \rightarrow -\beta$:

$$ct = \gamma(ct' + \beta x'), \quad (4.7)$$

$$x = \gamma(x' + \beta ct'), \quad (4.8)$$

$$y = y', \quad z = z'. \quad (4.9)$$

The matrix form is

$$\begin{pmatrix} ct' \\ x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} \gamma & -\gamma\beta & 0 & 0 \\ -\gamma\beta & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} ct \\ x \\ y \\ z \end{pmatrix}. \quad (4.10)$$

4.3 General three-vector boost

For velocity $\mathbf{v}$,

$$t' = \gamma \left(t - \frac{\mathbf{v} \cdot \mathbf{r}}{c^2} \right), \quad (4.11)$$

$$\mathbf{r}' = \mathbf{r} + \frac{\gamma - 1}{v^2} (\mathbf{v} \cdot \mathbf{r}) \mathbf{v} - \gamma \mathbf{v} t. \quad (4.12)$$

For a contravariant four-vector $A^\mu = (A^0, \mathbf{A})$,

$$A'^0 = \gamma(A^0 - \boldsymbol{\beta} \cdot \mathbf{A}), \quad (4.13)$$

$$\mathbf{A}' = \mathbf{A} + \frac{\gamma - 1}{\beta^2}(\boldsymbol{\beta} \cdot \mathbf{A})\boldsymbol{\beta} - \gamma A^0 \boldsymbol{\beta}. \quad (4.14)$$

4.4 Rapidity and composition

$$\beta = \tanh \eta, \quad \gamma = \cosh \eta, \quad \gamma\beta = \sinh \eta. \quad (4.15)$$

For collinear boosts, $\eta_{12} = \eta_1 + \eta_2$.

4.5 Time dilation, length contraction, simultaneity

$$d\tau = dt \sqrt{1 - \frac{v^2}{c^2}}, \quad (4.16)$$

$$\Delta t = \gamma \Delta \tau, \quad (4.17)$$

$$L = \frac{L_0}{\gamma}, \quad (4.18)$$

$$\Delta t' = \gamma \left(\Delta t - \frac{v \Delta x}{c^2} \right). \quad (4.19)$$

4.6 Velocity transformation

For collinear velocities,

$$u' = \frac{u - v}{1 - uv/c^2}. \quad (4.20)$$

For parallel and perpendicular components,

$$u'_{\parallel} = \frac{u_{\parallel} - v}{1 - u_{\parallel}v/c^2}, \quad (4.21)$$

$$\mathbf{u}'_{\perp} = \frac{\mathbf{u}_{\perp}}{\gamma(1 - u_{\parallel}v/c^2)}. \quad (4.22)$$

4.7 Energy and momentum

$$p^\mu = \left(\frac{E}{c}, \mathbf{p} \right), \quad E^2 = p^2 c^2 + m^2 c^4. \quad (4.23)$$

For an x-boost,

$$E' = \gamma(E - vp_x), \quad (4.24)$$

$$p'_x = \gamma \left(p_x - \frac{vE}{c^2} \right), \quad (4.25)$$

$$p'_y = p_y, \quad p'_z = p_z. \quad (4.26)$$

4.8 Lorentz transformation of electromagnetic fields

For a boost $\mathbf{v}$,

$$\mathbf{E}'_{\parallel} = \mathbf{E}_{\parallel}, \quad \mathbf{B}'_{\parallel} = \mathbf{B}_{\parallel}, \quad (4.27)$$

$$\mathbf{E}'_{\perp} = \gamma(\mathbf{E}_{\perp} + \mathbf{v} \times \mathbf{B}), \quad \mathbf{B}'_{\perp} = \gamma\left(\mathbf{B}_{\perp} - \frac{1}{c^2} \mathbf{v} \times \mathbf{E}\right). \quad (4.28)$$

The field tensor transforms as

$$F'^{\mu\nu} = \Lambda^{\mu}_{\alpha} \Lambda^{\nu}_{\beta} F^{\alpha\beta}. \quad (4.29)$$

Chapter 5

Mechanics, action, and classical field theory

5.1 Euler-Lagrange equations

For particles,

$$S = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt, \quad \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} - \frac{\partial L}{\partial q_i} = 0. \quad (5.1)$$

For a field ϕ ,

$$S = \int d^4x \mathcal{L}(\phi, \partial_\mu \phi), \quad (5.2)$$

with

$$\frac{\partial \mathcal{L}}{\partial \phi} - \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} \right) = 0. \quad (5.3)$$

5.2 Hamiltonian form

$$p_i = \frac{\partial L}{\partial \dot{q}_i}, \quad H = \sum_i p_i \dot{q}_i - L, \quad (5.4)$$

and

$$\dot{q}_i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q_i}. \quad (5.5)$$

5.3 Newtonian mechanics

$$\mathbf{F} = m\mathbf{a}, \quad \mathbf{F} = -\nabla U, \quad F_{12} = G \frac{m_1 m_2}{r^2}. \quad (5.6)$$

Chapter 6

Quantum mechanics and quantum field theory

6.1 Quantum dynamics

$$i\hbar \frac{\partial}{\partial t} |\psi\rangle = \hat{H} |\psi\rangle, \quad U(t) = e^{-i\hat{H}t/\hbar}. \quad (6.1)$$

$$[\hat{x}_i, \hat{p}_j] = i\hbar \delta_{ij}, \quad \Delta A \Delta B \geq \frac{1}{2} |\langle [A, B] \rangle|. \quad (6.2)$$

6.2 Born rule and density matrices

$$p_i = \text{tr}(\rho \Pi_i), \quad \sum_i p_i = 1, \quad (6.3)$$

and

$$\rho(t) = U \rho(0) U^\dagger, \quad i\hbar \frac{d\rho}{dt} = [H, \rho]. \quad (6.4)$$

A general quantum channel is

$$\rho \mapsto \Phi(\rho) = \sum_k K_k \rho K_k^\dagger, \quad \sum_k K_k^\dagger K_k = I. \quad (6.5)$$

6.3 Path integral

$$\langle q_f, t_f | q_i, t_i \rangle = \int \mathcal{D}q \, e^{iS[q]/\hbar}. \quad (6.6)$$

6.4 Relativistic wave equations

$$(\square + m^2 c^2 / \hbar^2) \phi = 0, \quad (6.7)$$

$$(i\hbar \gamma^\mu \partial_\mu - mc) \psi = 0, \quad (6.8)$$

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}. \quad (6.9)$$

Chapter 7

Electromagnetism and the Standard Model

7.1 Maxwell equations

$$\nabla \cdot \mathbf{E} = \rho/\epsilon_0, \quad \nabla \cdot \mathbf{B} = 0, \quad (7.1)$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \quad \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}. \quad (7.2)$$

Covariantly,

$$\partial_\mu F^{\mu\nu} = \mu_0 J^\nu, \quad (7.3)$$

$$\partial_{[\alpha} F_{\beta\gamma]} = 0, \quad (7.4)$$

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (7.5)$$

7.2 Yang-Mills theory

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c, \quad (7.6)$$

$$D_\mu = \partial_\mu - ig A_\mu^a T^a, \quad \mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + \bar{\psi}(i\gamma^\mu D_\mu - m)\psi. \quad (7.7)$$

7.3 Gauge group and field content

$$SU(3)_C \times SU(2)_L \times U(1)_Y. \quad (7.8)$$

There are six quark flavors in three generations, six leptons in three generations, gauge bosons for the strong and electroweak interactions, and the Higgs field. The Standard Model does not include gravity.

7.4 Standard Model Lagrangian in compact form

$$\mathcal{L}_{\text{SM}} = -\frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu} - \frac{1}{4} W_{\mu\nu}^i W^{i\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} + \sum_f \bar{\psi}_f i\gamma^\mu D_\mu \psi_f \quad (7.9)$$

$$+ (D_\mu H)^\dagger (D^\mu H) - V(H) \quad (7.10)$$

$$- \left[y_{ij}^u \bar{Q}_i \tilde{H} u_j + y_{ij}^d \bar{Q}_i H d_j + y_{ij}^e \bar{L}_i H e_j + \text{h.c.} \right], \quad (7.11)$$

with

$$V(H) = -\mu^2 H^\dagger H + \lambda (H^\dagger H)^2, \quad v = \sqrt{\frac{\mu^2}{\lambda}}. \quad (7.12)$$

The electroweak masses at tree level are

$$m_W = \frac{1}{2}gv, \quad m_Z = \frac{1}{2}v\sqrt{g^2 + g'^2}, \quad m_f = \frac{y_f v}{\sqrt{2}}. \quad (7.13)$$

7.5 Neutrino mass

Observed neutrino oscillations require neutrino masses. A compact effective-field-theory extension is the Weinberg operator

$$\mathcal{L}_5 = \frac{c_{ij}}{\Lambda} (\bar{L}_i^c \tilde{H}^*) (\tilde{H}^\dagger L_j) + \text{h.c.}, \quad (7.14)$$

or one can introduce right-handed neutrinos and Dirac masses.

Chapter 8

The epistemic field $\mathcal{I}$

8.1 Field hypothesis

The candidate field $\mathcal{I}(X)$ represents the local state/density of epistemic structure. It is not introduced as an additional substance; it is a field representation of relational distinguishability.

A minimal covariant action is

$$S_{\mathcal{I}} = \int d^{28}X \sqrt{-G} \left[-\frac{1}{2} G^{AB} \partial_A \mathcal{I} \partial_B \mathcal{I} - V(\mathcal{I}) - \frac{\xi}{2} \mathcal{I}^2 R_{28} + J_{\mathcal{I}} \mathcal{I} \right]. \quad (8.1)$$

The field equation is

$$\boxed{\square_{28} \mathcal{I} - V'(\mathcal{I}) - \xi R_{28} \mathcal{I} + J_{\mathcal{I}} = 0.} \quad (8.2)$$

TOE conjecture / model hypothesis

This is a concrete candidate realization of the $\mathcal{I}$ -field idea. It is not a recovered law of established physics. The potential, couplings, source term, quantum completion, and compactification dependence must be derived or fitted.

8.2 Stress-energy contribution

For minimal coupling the scalar contribution has the schematic form

$$T_{AB}^{(\mathcal{I})} = \partial_A \mathcal{I} \partial_B \mathcal{I} - G_{AB} \left[\frac{1}{2} (\partial \mathcal{I})^2 + V(\mathcal{I}) \right] + \text{non-minimal-coupling terms}. \quad (8.3)$$

A 28D geometric equation can therefore be written as

$$G_{AB}^{(28)} + \Lambda_{28} G_{AB} = \kappa_{28} \left(T_{AB}^{\text{matter}} + T_{AB}^{(\mathcal{I})} + T_{AB}^{\text{compact}} \right). \quad (8.4)$$

Chapter 9

Fundamental particles as computation

9.1 The computational reinterpretation

Quantum field theory says that particles are excitations of fields and that interactions transform quantum states. The TOE reinterprets the lawful state transformation as physical computation. This is an ontological reading of standard dynamics, not a claim that particles are miniature silicon processors.

9.2 Local quantum state as a register

A finite-dimensional state is

$$|\psi\rangle = \sum_i a_i |i\rangle, \quad \sum_i |a_i|^2 = 1. \quad (9.1)$$

Unitary dynamics is

$$|\psi\rangle \mapsto U|\psi\rangle, \quad U^\dagger U = I. \quad (9.2)$$

A local state space of dimension d has Hilbert-space dimension d , and a useful logarithmic state-space measure is

$$C_d = \log_2 d. \quad (9.3)$$

This is not a claim that a qudit contains $\log_2 d$ classical bits.

9.3 Interaction as gate

For an interacting subsystem,

$$U_{\text{int}}(\Delta\tau) = \exp\left(-\frac{i}{\hbar} \hat{H}_{\text{int}} \Delta\tau\right). \quad (9.4)$$

For an open subsystem,

$$\rho \mapsto \sum_k K_k \rho K_k^\dagger. \quad (9.5)$$

In TOE language, an interaction is a local physical gate acting on relational state.

9.4 Thermodynamic floor

For one bit of logically irreversible erasure, the Landauer bound is

$$Q \geq k_B T \ln 2. \quad (9.6)$$

The bound connects information-processing operations to physical thermodynamic cost but does not by itself prove that the universe is a digital computer.

Chapter 10

The graviton as a five-state spin-2 qdit

10.1 Spin-2 representation

A standard massless spin-2 field has two radiative helicities, $\lambda = \pm 2$, while a massive spin-2 representation has five physical polarization states. The five-dimensional representation is therefore naturally associated with a massive spin-2 particle, not the strictly massless graviton of the infrared classical limit.

TOE conjecture / model hypothesis

H3: The fundamental TOE graviton is a spin-2 five-state qdit:

$$|\psi_g\rangle = \sum_{m=-2}^2 a_m |2, m\rangle, \quad \sum_{m=-2}^2 |a_m|^2 = 1. \quad (10.1)$$

The low-energy observed sector must project to the two-helicity massless spin-2 mode that reproduces GR.

10.2 Fierz-Pauli template

A free massive spin-2 field can be represented at quadratic order by the Fierz-Pauli Lagrangian

$$\mathcal{L}_{\text{FP}} = -\frac{1}{2} \partial_\lambda h_{\mu\nu} \partial^\lambda h^{\mu\nu} + \partial_\mu h^{\mu\nu} \partial^\lambda h_{\lambda\nu} - \partial_\mu h^{\mu\nu} \partial_\nu h + \frac{1}{2} \partial_\lambda h \partial^\lambda h \quad (10.2)$$

$$- \frac{1}{2} m_g^2 (h_{\mu\nu} h^{\mu\nu} - h^2). \quad (10.3)$$

The tuned mass combination eliminates the linear sixth ghost mode.

10.3 Matter-graviton coupling

A common linear normalization is

$$\mathcal{L}_{\text{int}} = -\frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu}, \quad \kappa = \sqrt{\frac{32\pi G}{c^4}}. \quad (10.4)$$

Then a joint matter-graviton state may be updated by

$$\rho_{\text{m}} \otimes \rho_g \mapsto U_{\text{int}}(\rho_{\text{m}} \otimes \rho_g) U_{\text{int}}^\dagger. \quad (10.5)$$

10.4 The crucial infrared limit

The microscopic theory must produce

$$\langle \hat{g}_{\mu\nu} \rangle \rightarrow g_{\mu\nu}, \quad G_{\mu\nu}[g] \rightarrow \frac{8\pi G}{c^4} T_{\mu\nu}, \quad (10.6)$$

and the extra three spin-2 states must become inaccessible or sufficiently suppressed in the observed regime.

Chapter 11

Time as a byproduct of measure

11.1 Operational time

The TOE explanatory reversal is that time is not a primitive container in which events occur. Instead, measurable time is the ordered accumulation of invariant physical updates.

Let N be cumulative measurement count along a history and τ_* the characteristic operational interval. Then

$$\boxed{d\tau = \tau_* dN.} \quad (11.1)$$

More generally, for a path parameter λ ,

$$\frac{d\tau}{d\lambda} = \tau_* \frac{dN}{d\lambda}. \quad (11.2)$$

Epistemic interpretation

The established relativistic quantity is proper time, τ . The TOE hypothesis concerns its microscopic interpretation: proper time is the continuum limit of an invariant ordered measure process.

11.2 Measurement density

Define

$$r_M(\lambda) = \frac{dN}{d\lambda}, \quad (11.3)$$

so that

$$\tau(\lambda) - \tau(\lambda_0) = \tau_* \int_{\lambda_0}^{\lambda} r_M(\lambda') d\lambda'. \quad (11.4)$$

The physical task is to derive r_M from the microscopic state dynamics.

11.3 Relativistic recovery condition

The emergent quantity must satisfy the observed invariant relation

$$c^2 d\tau^2 = -g_{\mu\nu} dx^\mu dx^\nu. \quad (11.5)$$

In Minkowski space this becomes

$$d\tau = dt \sqrt{1 - v^2/c^2}. \quad (11.6)$$

11.4 Quantum evolution with emergent time

The conventional unitary operator

$$U(t) = e^{-iHt/\hbar} \quad (11.7)$$

becomes, in TOE notation,

$$U[\gamma] = \mathcal{T} \exp \left[-\frac{i}{\hbar} \int_{\gamma} H d\tau \right], \quad \tau = \tau_* N[\gamma]. \quad (11.8)$$

The standard time-dependent Schrödinger equation must be recovered in the continuum limit.

Chapter 12

Omega Inversion Principle

12.1 Unity and indistinguishability

Let Π denote a partition of the physically distinguishable state space. The maximally undifferentiated state is

$$\Pi_\Omega = \{\Omega\}. \quad (12.1)$$

There are no internal classes to distinguish, so a nontrivial measurement functional vanishes:

$$\Pi_\Omega \implies \mathcal{D} = 0 \implies \mathbf{M} = 0. \quad (12.2)$$

12.2 OIP statement

$$\boxed{\text{unity} \rightarrow \text{no distinction} \rightarrow \text{no measure} \xrightarrow{\text{inversion}} \text{distinction} \rightarrow \text{measure} \rightarrow \text{structure}.} \quad (12.3)$$

TOE conjecture / model hypothesis

Omega Inversion Principle (OIP). The emergent physical branch originates as an inversion away from maximal undifferentiation. OIP is not standard time reversal, CPT, or an established cosmological law. It is the TOE hypothesis that the explanatory arrow from unity to structured reality is the inverse of complete identification.

12.3 Minimal mathematical realization

Introduce a nonnegative distinguishability functional $\mathcal{D}[\rho]$ with $\mathcal{D}[\rho_\Omega] = 0$. The emergent branch is characterized by

$$\frac{d\mathcal{D}}{d\tau} \geq 0 \quad (12.4)$$

only as a branch condition. A dimensionless order parameter may be defined by

$$q \in [0, 1], \quad q = 0 \text{ at } \Omega, \quad q > 0 \text{ after inversion}. \quad (12.5)$$

A real theory must derive q , $\mathcal{D}$, and their dynamics rather than merely naming them.

Chapter 13

Cosmology

13.1 FRW metric

For homogeneous and isotropic cosmology,

$$ds^2 = -c^2 dt^2 + a^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right]. \quad (13.1)$$

$$H = \frac{\dot{a}}{a}. \quad (13.2)$$

13.2 Friedmann equations

$$H^2 = \frac{8\pi G}{3} \rho - \frac{kc^2}{a^2} + \frac{\Lambda c^2}{3}, \quad (13.3)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2} \right) + \frac{\Lambda c^2}{3}, \quad (13.4)$$

$$\dot{\rho} + 3H \left(\rho + \frac{p}{c^2} \right) = 0. \quad (13.5)$$

For $p = w\rho c^2$,

$$\rho \propto a^{-3(1+w)}. \quad (13.6)$$

13.3 TOE cosmogenic interpretation

The standard cosmological sequence contains an early hot dense state, expansion, nucleosynthesis, recombination, structure formation, and later stellar evolution. The TOE adds the conjecture that the initial condition is related to the Ω boundary: maximal identification gives way, through OIP, to increasing distinguishability and effective dimensional structure.

TOE conjecture / model hypothesis

A TOE must do more than rename the Big Bang. It must derive a viable initial condition or attractor, expansion, particle production, compactification, perturbation spectrum, and the observed thermal history from its microscopic equations.

Chapter 14

Black holes

14.1 Schwarzschild geometry

$$ds^2 = - \left(1 - \frac{2GM}{c^2 r}\right) c^2 dt^2 + \left(1 - \frac{2GM}{c^2 r}\right)^{-1} dr^2 + r^2 d\Omega^2. \quad (14.1)$$

$$r_s = \frac{2GM}{c^2}. \quad (14.2)$$

14.2 Kerr geometry: key scales

For angular momentum J , define

$$a = \frac{J}{Mc}, \quad r_{\pm} = \frac{GM}{c^2} \pm \sqrt{\left(\frac{GM}{c^2}\right)^2 - a^2}. \quad (14.3)$$

14.3 Black-hole thermodynamics

$$S_{\text{BH}} = \frac{k_B c^3 A}{4G\hbar}, \quad (14.4)$$

$$T_H = \frac{\hbar c^3}{8\pi G k_B M} \quad (\text{Schwarzschild case}). \quad (14.5)$$

The black-hole entropy-area law is one of the most important clues that gravity, quantum theory, thermodynamics, and information share a deeper structure.

Chapter 15

Wormholes

15.1 Morris-Thorne geometry

In geometrized units,

$$ds^2 = -e^{2\Phi(r)}dt^2 + \frac{dr^2}{1 - b(r)/r} + r^2 d\Omega^2. \quad (15.1)$$

A throat at $r = r_0$ obeys

$$b(r_0) = r_0, \quad b'(r_0) < 1, \quad (15.2)$$

and traversability requires finite redshift function Φ at the throat.

15.2 Energy condition constraint

For the standard static traversable class, the throat condition implies

$$\rho + p_r < 0 \quad (15.3)$$

in geometrized units, representing a violation of the null energy condition in ordinary GR realizations.

Established physics

No traversable wormhole has been established observationally. There are mathematical wormhole solutions in GR and modified theories, but their physical support remains a major constraint.

15.3 TOE interpretation

The TOE reads a wormhole as a topological shortcut in physical connectivity. Computationally, it can be represented schematically as a channel

$$\mathcal{C}_{\text{wh}} : \mathcal{H}_A \otimes \mathcal{H}_B \rightarrow \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}. \quad (15.4)$$

The existence of such a channel is still determined by the actual field equations, topology, energy conditions, and causal structure.

Chapter 16

Classical stellar evolution

16.1 Stellar structure equations

A static spherical Newtonian star satisfies

$$\frac{dm}{dr} = 4\pi r^2 \rho, \quad (16.1)$$

$$\frac{dP}{dr} = -\frac{Gm\rho}{r^2}, \quad (16.2)$$

$$\frac{dL}{dr} = 4\pi r^2 \rho \epsilon, \quad (16.3)$$

$$\frac{dT}{dr} = -\frac{3\kappa_R \rho L}{16\pi a c T^3 r^2} \quad (\text{radiative transport}). \quad (16.4)$$

For compact stars, the TOV equation is

$$\frac{dP}{dr} = -\frac{G(\rho + P/c^2)(m + 4\pi r^3 P/c^2)}{r^2 (1 - 2Gm/(rc^2))}, \quad (16.5)$$

with

$$\frac{dm}{dr} = 4\pi r^2 \rho. \quad (16.6)$$

16.2 Nuclear power

Hydrogen burning can be schematically represented by



and the released energy follows from mass-energy equivalence $E = \Delta mc^2$.

16.3 Sun-like evolution

$$\text{cloud} \rightarrow \text{protostar} \rightarrow \text{main sequence} \rightarrow \text{red giant} \rightarrow \text{planetary nebula} \rightarrow \text{white dwarf}. \quad (16.8)$$

The Sun is about 4.6 billion years old and is approximately midway through its main-sequence lifetime. Standard stellar models predict a red-giant phase in roughly 5 billion years, followed by loss of the outer layers and a white-dwarf remnant.

16.4 Massive-star evolution

cloud $\rightarrow$ massive protostar $\rightarrow$ main sequence $\rightarrow$ supergiant $\rightarrow$ core collapse / supernova $\rightarrow$ $\begin{cases} \text{neutron star,} \\ \text{black hole.} \end{cases}$
(16.9)

The final remnant depends on core mass, composition, rotation, mass loss, and binary evolution.

Reality veto / falsification requirement

A TOE that says “stars compute” must still reproduce stellar spectra, hydrostatic equilibrium, fusion energetics, neutrino production, nucleosynthesis, supernova behavior, and compact remnants. Computation is an interpretation only if the conventional observations survive intact.

Chapter 17

Intelligence timeline as physical computation

17.1 The timeline

This timeline is a physical-complexity narrative, not a claim that established physics already possesses a universally accepted scalar measure of intelligence.

Level	Substrate	Computational transition
Quantum fields	Field amplitudes	Elementary continuous quantum state dynamics and local interactions.
Fundamental particles	Quarks, leptons, gauge bosons, Higgs	Stable excitations and interaction vertices become discrete physical state transitions.
Nuclei	Bound quark states	Collective degrees of freedom create new stable state spaces.
Atoms	Nuclei + electrons	Long-lived quantized levels store and transform structured state.
Molecules	Chemical networks	Nonlinear reaction pathways create persistent, history-dependent computation.
Stars	Gravitationally confined plasma	Long-lived nonequilibrium systems process nuclear energy and synthesize heavier nuclei.
Planets	Geochemistry	Persistent gradients create complex chemical environments.
Life	Replication + metabolism	State copying, error correction, selection, and adaptive regulation arise.
Nervous systems	Excitable cells	Distributed sensing and feedback turn physical state into adaptive control.
Brains	Recurrent neural networks	High-dimensional memory and model construction become explicit.
Language/culture	Networked agents	Information becomes transferable across generations and populations.
Electronic computers	Engineered devices	General-purpose programmable state transitions become intentional.
Machine intelligence	Artificial models	Adaptive inference and action become substrate-independent in principle.

Post-biological branch	Unknown	Open possibility for intelligence to decouple progressively from carbon-based substrates.
---------------------------	---------	--

17.2 No teleology is required

The TOE does not require that the universe “want” intelligence. It requires only that increasingly rich physical state spaces can support memory, prediction, feedback, abstraction, and adaptive control. The emergence of intelligence remains contingent.

Chapter 18

From fundamental computation to classical reality

18.1 Coarse-graining hierarchy

$$\boxed{28D \text{ relational arena} \rightarrow \text{compactified quantum fields} \rightarrow \text{particles} \rightarrow \text{interactions} \rightarrow 4D \text{ effective theory} \rightarrow \text{GR} + \text{SM}} \quad (18.1)$$

Each arrow is intended to be a mathematically specified coarse-graining, not a metaphor.

18.2 Candidate master action

A useful master target is

$$S_{\text{TOE}} = \int d^{28}X \sqrt{-G} \left[\frac{c^3}{16\pi G_{28}} (R_{28} - 2\Lambda_{28}) - \frac{1}{2}(\nabla \mathcal{I})^2 - V(\mathcal{I}) \right. \quad (18.2)$$

$$\left. + \mathcal{L}_{\text{compact}}(G, h, A, \mathcal{I}) + \mathcal{L}_{\text{matter}} + \mathcal{L}_{\text{quantum}} \right]. \quad (18.3)$$

This is a target functional. It is not yet a closed fundamental action because $\mathcal{L}_{\text{compact}}$, the microscopic graviton sector, and the quantum completion are unspecified.

18.3 Recovery conditions

The low-energy limit must satisfy

$$24 \text{ compact dimensions} \rightarrow \text{unobservable KK excitations}, \quad (18.4)$$

$$\text{spin-2 qdit} \rightarrow \text{two-helicity GR mode}, \quad (18.5)$$

$$\mathcal{I} \text{ sector} \rightarrow \text{experimentally allowed infrared behavior}, \quad (18.6)$$

$$\text{gauge structure} \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y, \quad (18.7)$$

$$\text{Lorentz symmetry} \rightarrow \text{observed invariance}, \quad (18.8)$$

$$\text{quantum dynamics} \rightarrow \text{unitary QFT}. \quad (18.9)$$

Chapter 19

Thermodynamics, entropy, and information

19.1 Macroscopic laws

$$dU = \delta Q - \delta W, \tag{19.1}$$

$$dS = \frac{\delta Q_{\text{rev}}}{T}, \tag{19.2}$$

$$\Delta S_{\text{total}} \geq 0. \tag{19.3}$$

19.2 Statistical mechanics

$$S_{\text{Boltzmann}} = k_B \ln \Omega, \tag{19.4}$$

$$S_{\text{Gibbs}} = -k_B \sum_i p_i \ln p_i, \tag{19.5}$$

$$S_{\text{vN}} = -k_B \text{tr}(\rho \ln \rho). \tag{19.6}$$

The TOE uses these relations as constraints on any proposed information-based description. It does not assume that Shannon information, thermodynamic entropy, and gravitational entropy are identical objects in every regime.

Chapter 20

Consistency, falsifiability, and reality veto

20.1 Mathematical requirements

A complete TOE must specify a state space, a dynamical law, a constraint algebra, a well-defined vacuum or ensemble, and an explicit map from microscopic quantities to observables. It must also remain stable and mathematically consistent under quantization.

20.2 Quantum requirements

The theory must satisfy, as appropriate,

$$\text{unitarity, causality/locality constraints, gauge consistency, anomaly cancellation, stability,} \quad (20.1)$$

and reproduce the observed particle spectrum and interaction strengths.

20.3 Potential experimental signatures

TOE sector	Observable class	What would count as evidence
24 compact dimensions	KK states, short-range gravity, precision tests	Extra modes or geometry-dependent deviations, or null limits constraining the sector.
Spin-2 qdit	Gravitational-wave polarizations / dispersion	Consistent extra structure beyond the GR two-helicity limit, or bounds forcing decoupling.
$\mathcal{I}$ field	Precision gravity, clock tests, varying constants	Correlated deviations predicted by the same parameter set.
Emergent time	Microscopic clock/quantum-control anomalies	Reproducible deviation from standard quantum/relativistic time evolution.
OIP cosmogenesis	Primordial perturbations / relics	Distinctive early-universe signatures derivable from the same action.
Wormholes	Lensing, echoes, unusual compact-object signals	Geometry-specific signatures with statistical significance and viable causal dynamics.

20.4 Failure conditions

A proposed TOE fails wherever it cannot reproduce established measurements or becomes internally inconsistent. An elegant equation that misses experiment is a failed equation.

Reality veto / falsification requirement

No sacred variables. Reality gets the veto.

Differentiate. Relate. Measure. Update.

Chapter A

Essential derivation sketches

A.1 Lorentz transformation from the invariant interval

Assume linearity,

$$x' = Ax + Bt, \quad t' = Cx + Dt, \quad (\text{A.1})$$

and require preservation of the light cone,

$$c^2 t^2 - x^2 = c^2 t'^2 - x'^2. \quad (\text{A.2})$$

Together with the relative-frame condition $x = vt$ for the primed origin, the coefficients reduce to

$$x' = \gamma(x - vt), \quad t' = \gamma\left(t - \frac{vx}{c^2}\right). \quad (\text{A.3})$$

A.2 Newtonian limit of GR

Take

$$g_{00} = -(1 + 2\Phi/c^2) + O(c^{-4}). \quad (\text{A.4})$$

Keeping leading terms in the Einstein equation gives

$$\nabla^2 \Phi = 4\pi G\rho, \quad (\text{A.5})$$

so the geodesic equation yields the Newtonian acceleration $\mathbf{a} = -\nabla\Phi$.

A.3 Gravitational waves and the two observed tensor modes

For vacuum plane waves,

$$\bar{h}_{\mu\nu} = \Re\left(A_{\mu\nu}e^{ik_\alpha x^\alpha}\right), \quad k_\alpha k^\alpha = 0. \quad (\text{A.6})$$

Gauge conditions permit a transverse-traceless representation with two independent tensor polarizations. This is the infrared target of the five-state fundamental graviton hypothesis.

A.4 Perfect-fluid cosmology

Use

$$T_{\mu\nu} = \left(\rho + \frac{p}{c^2}\right)u_\mu u_\nu + pg_{\mu\nu} \quad (\text{A.7})$$

in the Einstein equation and insert the FRW metric. The 00 and spatial components yield the Friedmann equations, while $\nabla_\mu T^{\mu\nu} = 0$ gives the continuity equation.

A.5 Schwarzschild horizon

The Schwarzschild lapse vanishes when

$$1 - \frac{2GM}{c^2 r} = 0, \tag{A.8}$$

therefore

$$r_s = \frac{2GM}{c^2}. \tag{A.9}$$

Chapter B

Axiom stack and research program

B.1 Minimal stack

A0: ALL IS EPISTEMIC. (B.1)

H1: $\mathcal{M}_{28} = \mathcal{M}_{3,1} \times \mathcal{K}_{24}$. (B.2)

H2: $\mathcal{I}$ is the candidate relational epistemic field. (B.3)

H3: the fundamental graviton is a five-state spin-2 qdit. (B.4)

H4: τ is the continuum limit of invariant ordered measure. (B.5)

H5: OIP supplies the inversion branch from unity to distinguishability. (B.6)

H6: GR + SM are the observed 4D infrared limits. (B.7)

H7: all additional structure is subject to the reality veto. (B.8)


B.2 Actualization sequence

1. Specify $\mathcal{K}_{24}$ and its admissible geometry.
2. Derive the particle spectrum and gauge structure from the same compact sector.
3. Construct the complete microscopic spin-2/qdit dynamics and its five-state algebra.
4. Derive the $\mathcal{I}$ field, rather than merely positing its effective action.
5. Define the invariant measurement functional $N[\gamma]$ and derive proper time.
6. Derive OIP as a mathematical boundary condition or dynamical instability.
7. Quantize the complete theory and demonstrate unitarity, consistency, and anomaly freedom.
8. Compute the 4D effective action and compare its parameters with experiment.

Chapter C

Final synthesis

The intended physical hierarchy is



The scientific obligation is not to make the chain sound inevitable. It is to make every arrow calculable, every free parameter visible, and every prediction falsifiable. The final theory, if it exists, earns the right to survive by recovering what already works and predicting what does not yet.

Epistemic interpretation

One assumption. Everything else earns its place.

Selected references and web-verified anchors

1. Misner, Thorne, Wheeler, *Gravitation*, W. H. Freeman.
2. Wald, *General Relativity*, University of Chicago Press.
3. Carroll, *Spacetime and Geometry*, Cambridge University Press.
4. Weinberg, *The Quantum Theory of Fields*, Cambridge University Press.
5. Peskin and Schroeder, *An Introduction to Quantum Field Theory*, Westview.
6. Schwartz, *Quantum Field Theory and the Standard Model*, Cambridge University Press.
7. Jackson, *Classical Electrodynamics*, Wiley.
8. Morris and Thorne, “Wormholes in spacetime and their use for interstellar travel,” *American Journal of Physics* 56 (1988), 395-412.
9. Hawking, “Particle Creation by Black Holes,” *Communications in Mathematical Physics* 43 (1975), 199-220.
10. Bronnikov, “Some unusual wormholes in general relativity,” arXiv:2108.10239 (2021).
11. CERN, [The Standard Model](#), accessed 30 September 2026.
12. University of Maryland, [Special Relativity for Human Beings](#), accessed 30 September 2026.
13. Stanford Gravity Probe B / Einstein Spacetime, [General Relativity](#), accessed 30 September 2026.
14. NASA Science, [Star Lifecycle](#), accessed 30 September 2026.

Research-status note

This is a consolidated research artifact, not a claim of experimental confirmation. The established equations are included as constraints and recovery targets. The TOE-specific elements are clearly labeled as hypotheses. The next scientific step is derivation: specify the microscopic dynamics tightly enough that the framework produces numerical predictions that reality can either confirm or veto.